

POSSIBLE SIDE EFFECTS OF VACCINES

WHAT YOU NEED TO KNOW

Like any other drug, vaccines too may have **undesirable effects**, such as:

LOCAL UNDESIRABLE EFFECTS,
I.E., AT THE INJECTION SITE OR,
LESS COMMONLY,
GENERAL UNDESIRABLE EFFECTS, SUCH AS ALLERGIC REACTIONS!



SHOULDER AND LEG

Redness or swelling at the injection site: it does not depend on the healthcare worker and it goes away in a couple of weeks²



BODY

- Your child "is a little sluggish": s/he is not feeling well and is irritable for 24-48 hours but is easy to comfort²
- Skin rash (exanthema): may occur more often after the measles vaccine, 7-10 days after injection and goes away spontaneously³



ABDOMEN

Diarrhoea: may occur with oral vaccines such as the Rotavirus vaccine⁴



FEVER AND PAIN

- **Fever:** is generally between 38 and 38.5°C, does not last long (24-48 hours), and has no consequences
- **Pain:** at the injection site, regardless of redness or swelling



If necessary, use of **ibuprofen or paracetamol** is effective in treating both the fever and pain, thus helping your child to feel better²



scan the
QR code
for more
special
contents

DON'T FORGET THAT THE UNDESIRABLE EFFECTS OF VACCINES:

- are mostly unpredictable, random and mild (vaccines are safe!)⁴
- can be easily controlled, starting with the fever and pain²
- are always hugely outweighed by the benefits of vaccination compared to the risk of diseases⁴

